

April 16, 2003

IN THE SPECIFICATION

Please replace the paragraph beginning at page 10, line 27, with the following rewritten paragraph:

Figure 4: Cyclization using bi-modal peptides. (a) Deprotection and cyclisation. Synthesis of bi-modal peptides which have complimentary ligands at their N- and C- termini allows the cyclisation of these peptides in aqueous buffers (i) Ligation. (ii) Deprotection and ligation. (iii) Cleavage of the cyclic peptide from the base labile handle. Example; The peptides shown are from Table 1 and represent major protective epitopes on PrtR 27 or PrtK39 (SEQ ID NO:4 and SEQ ID NO:5). (a) Ligation. 95% aqueous TFA. Ligation can be monitored by reverse phase analytical HPLC and mass spectrometry. Ligation conditions can be varied to include scavengers commonly used in peptide synthesis and different acidic conditions to enhance the Friedal-Craft alkylation. (b) Deprotection and ligation. The S-acetyl protecting group can removed by aqueous hydroxylamine 0.05 M, pH 7.3. Ligation, 6 M aqueous guanidine hydrochloride and 0.05 M EDTA pH 6.4-6.5 adjusted by 1 M Tris.HCl under nitrogen. The ligation strategy can also be accomplished on the solid phase. By selecting which ligand to introduce at the N- and C- terminal parallel and anti-parallel cyclic peptides can be synthesised.

Please replace the paragraph beginning at page 11, line 8, with the following rewritten paragraph:

Figure 5: Synthesis of multivalent multiple antigenic peptides (MAPs) using alternate ligation chemistries (SEQ ID NO:4 and SEQ ID NO:5). By using different ligation strategies a

vareity of peptides can be ligated onto a single multiple antigenic peptide. The example shown is of peptides listed in Table 1. (a) Ligation, 95% aqueous TFA. Ligation can be monitored by reverse phase analytical HPLC and mass spectrometry. Deprotection, Alloc can removed by palladium(0)-catalyzed allyl group transfer to a basic receptor. after purification the second peptide can be ligated on to the MAP, (c) 8 M urea and 0.1 M NaH₂PO₄ (pH range 3-4.7).

Please replace the paragraph beginning at page 11, line 18, with the following rewritten paragraph:

Figure 6: Serum IgG antibody responses assessed by ELISA to *Porphyromonas gingivalis* PrtR-27 overlapping peptides (SEQ ID NOs:13-24, 33, and 25-32, respectively, from origin). Twenty one PIN-bound peptides were probed with normal mouse sera (■), protective mouse sera (□), normal human serum (■), patient D24 sera (■), patient H10 sera (■) and patient D20 sera (■). ELISAs were developed as per Example 1.

Substitute the Sequence Listing submitted herewith for the Sequence Listing filed October 2, 2001.

O'BRIEN-SIMPSON

Serial No. 09/914,612

April 16, 2003

IN THE DRAWINGS

The attached replacement sheets of drawings include changes to Figs. 4, 5 and 6. The replacement sheets, Figs. 1-7, replace the original sheets including Figs. 4, 5 and 6 filed August 31, 2001.

Attachment: Replacement Sheets (Figs. 1-7)